#### Amendments to the Claims

This listing of claims will replace all prior versions, and listings of claims in the application.

 (Currently amended) A composition comprising a synergistically effective active compound combination amount of anthranilamides an anthranilamide of the formula (I)

in which

A<sup>1</sup> and A<sup>2</sup> independently of one another represent oxygen or sulfur,

X<sup>1</sup> represents N or CR<sup>10</sup>,

R<sup>1</sup> represents hydrogen or represents in each case optionally mono- or polysubstituted C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>2</sub>-C<sub>6</sub>-alkenyl, C<sub>2</sub>-C<sub>6</sub>-alkynyl or C<sub>3</sub>-C<sub>6</sub>-cycloalkyl, where the substituents independently of one another may be selected from the group consisting of R<sup>6</sup>, halogen, cyano, nitro, hydroxyl, C<sub>1</sub>-C<sub>4</sub>-alkoxy, C<sub>1</sub>-C<sub>4</sub>-alkylthio, C<sub>1</sub>-C<sub>4</sub>-alkylsulfinyl, C<sub>1</sub>-C<sub>4</sub>-alkylsulfonyl, C<sub>2</sub>-C<sub>4</sub>-alkoxycarbonyl, C<sub>1</sub>-C<sub>4</sub>-alkylamino, C<sub>2</sub>-C<sub>8</sub>-dialkylamino, C<sub>3</sub>-C<sub>6</sub>-cycloalkylamino, (C<sub>1</sub>-C<sub>4</sub>-alkyl)C<sub>3</sub>-C<sub>6</sub>-cycloalkylamino and R<sup>11</sup>,

R<sup>2</sup> represents hydrogen, C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>2</sub>-C<sub>6</sub>-alkenyl, C<sub>2</sub>-C<sub>6</sub>-alkynyl, C<sub>3</sub>-C<sub>6</sub>-cycloalkyl, C<sub>1</sub>-C<sub>4</sub>-alkoxy, C<sub>1</sub>-C<sub>4</sub>-alkylamino, C<sub>2</sub>-C<sub>8</sub>-dialkylamino, C<sub>3</sub>-C<sub>6</sub>-cycloalkylamino, C<sub>2</sub>-C<sub>6</sub>-alkoxycarbonyl or C<sub>2</sub>-C<sub>6</sub>-alkylcarbonyl,

R<sup>3</sup> represents hydrogen, R<sup>11</sup> or represents in each case optionally mono- or polysubstituted C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>2</sub>-C<sub>6</sub>-alkenyl, C<sub>2</sub>-C<sub>6</sub>-alkynyl, C<sub>3</sub>-C<sub>6</sub>-cycloalkyl, where the substituents independently of one another may be

selected from the group consisting of R<sup>6</sup>, halogen, cyano, nitro, hydroxyl, C<sub>1</sub>-C<sub>4</sub>-alkoxy, C<sub>1</sub>-C<sub>4</sub>-haloalkoxy, C<sub>1</sub>-C<sub>4</sub>-alkylthio, C<sub>1</sub>-C<sub>4</sub>-alkylsulfinyl, C<sub>1</sub>-C<sub>4</sub>-alkylsulfonyl, C<sub>2</sub>-C<sub>6</sub>-alkoxycarbonyl, C<sub>2</sub>-C<sub>6</sub>-alkylcarbonyl, C<sub>3</sub>-C<sub>6</sub>-trialkylsilyl, R<sup>11</sup>, phenyl, phenoxy and a 5- or 6-membered heteroaromatic ring, where each phenyl, phenoxy and 5- or 6-membered heteroaromatic ring may optionally be substituted and where the substituents independently of one another may be selected from one to three radicals W or one or more radicals R<sup>12</sup>, or

R<sup>2</sup> and R<sup>3</sup> may be attached to one another and form the ring M,

 $R^4$ represents hydrogen, C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>2</sub>-C<sub>6</sub>-alkenyl, C<sub>2</sub>-C<sub>6</sub>-alkynyl, C<sub>3</sub>-C<sub>6</sub>cycloalkyl, C<sub>1</sub>-C<sub>6</sub>-haloalkyl, C<sub>2</sub>-C<sub>6</sub>-haloalkenyl, C<sub>2</sub>-C<sub>6</sub>-haloalkynyl, C<sub>3</sub>-C<sub>6</sub>-halocycloalkyl, halogen, cyano, nitro, hydroxyl, C<sub>1</sub>-C<sub>4</sub>-alkoxy, C<sub>1</sub>-C<sub>4</sub>haloalkoxy, C<sub>1</sub>-C<sub>4</sub>-alkylthio, C<sub>1</sub>-C<sub>4</sub>-alkylsulfinyl, C<sub>1</sub>-C<sub>4</sub>-alkylsulfonyl, C<sub>1</sub>-C<sub>4</sub>-haloalkylthio, C<sub>1</sub>-C<sub>4</sub>-haloalkylsulfinyl, C<sub>1</sub>-C<sub>4</sub>-haloalkylsulfonyl, C<sub>1</sub>-C<sub>4</sub>-C<sub>2</sub>-C<sub>8</sub>-dialkylamino, C<sub>3</sub>-C<sub>6</sub>-cycloalkylamino, alkylamino, trialkylsilyl or represents in each case optionally mono- or polysubstituted phenyl, benzyl or phenoxy, where the substituents independently of one another may be selected from the group consisting of C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>2</sub>-C<sub>4</sub>alkenyl, C<sub>2</sub>-C<sub>4</sub>-alkynyl, C<sub>3</sub>-C<sub>6</sub>-cycloalkyl, C<sub>1</sub>-C<sub>4</sub>-haloalkyl, C<sub>2</sub>-C<sub>4</sub>haloalkenyl, C2-C4-haloalkynyl, C3-C6-halocycloalkyl, halogen, cyano, nitro, C<sub>1</sub>-C<sub>4</sub>-alkoxy, C<sub>1</sub>-C<sub>4</sub>-haloalkoxy, C<sub>1</sub>-C<sub>4</sub>-alkylthio, C<sub>1</sub>-C<sub>4</sub>-alkylsulfinyl, C<sub>1</sub>-C<sub>4</sub>-alkylsulfonyl, C<sub>1</sub>-C<sub>4</sub>-alkylamino, C<sub>2</sub>-C<sub>8</sub>-dialkylamino, C<sub>3</sub>-C<sub>6</sub>-cycloalkylamino, C<sub>3</sub>-C<sub>6</sub>-(alkyl)cycloalkylamino, C<sub>2</sub>-C<sub>4</sub>-alkylcarbonyl, C<sub>2</sub>-C<sub>6</sub>-alkoxycarbonyl, C<sub>2</sub>-C<sub>6</sub>-alkyl-aminocarbonyl, C<sub>3</sub>-C<sub>8</sub>-dialkylaminocarbonyl and C<sub>3</sub>-C<sub>6</sub>-trialkylsilyl.

 $R^5$  and  $R^8$  in each case independently of one another represent hydrogen, halogen or represent in each case optionally substituted  $C_1$ - $C_4$ -alkyl,  $C_1$ - $C_4$ -haloalkyl,  $R^{12}$ , G, J, -OJ, -OG, -S(O)<sub>p</sub>-J, -S(O)<sub>p</sub>-G, -S(O)<sub>p</sub>-phenyl, where the substituents independently of one another may be selected from one to three radicals W or from the group consisting of  $R^{12}$ ,  $C_1$ - $C_{10}$ -alkyl,  $C_2$ - $C_6$ -alkynyl,  $C_1$ - $C_4$ -alkoxy and  $C_1$ - $C_4$ -alkythio, where each

substituent may be substituted by one or more substituents independently of one another selected from the group consisting of G, J,  $R^6$ , halogen, cyano, nitro, amino, hydroxyl,  $C_1$ - $C_4$ -alkoxy,  $C_1$ - $C_4$ -haloalkoxy,  $C_1$ - $C_4$ -alkylthio,  $C_1$ - $C_4$ -alkylsulfinyl,  $C_1$ - $C_4$ -alkyl-sulfonyl,  $C_1$ - $C_4$ -haloalkylsulfinyl,  $C_1$ - $C_4$ -haloalkylsulfonyl,  $C_1$ - $C_4$ -alkylamino,  $C_2$ - $C_8$ -dialkylamino,  $C_3$ - $C_6$ -trialkylsilyl, phenyl and phenoxy, where each phenyl or phenoxy ring may optionally be substituted and where the substituents independently of one another may be selected from one to three radicals W or one or more radicals  $R^{12}$ ,

- in each case independently of one another represent a 5- or 6-membered nonaromatic carbocyclic or heterocyclic ring which optionally contains one or two ring members from the group consisting of C(=O), SO and S(=O)<sub>2</sub> and which may optionally be substituted by one to four substituents independently of one another selected from the group consisting of C<sub>1</sub>-C<sub>2</sub>-alkyl, halogen, cyano, nitro and C<sub>1</sub>-C<sub>2</sub>-alkoxy, or independently of one another represent C<sub>2</sub>-C<sub>6</sub>-alkenyl, C<sub>2</sub>-C<sub>6</sub>-alkynyl, C<sub>3</sub>-C<sub>7</sub>-cycloalkyl, (cyano)C<sub>3</sub>-C<sub>7</sub>-cycloalkyl, (C<sub>1</sub>-C<sub>4</sub>-alkyl)C<sub>3</sub>-C<sub>6</sub>-cycloalkyl, (C<sub>3</sub>-C<sub>6</sub>-cycloalkyl)C<sub>1</sub>-C<sub>4</sub>-alkyl, where each cycloalkyl, (alkyl)cycloalkyl and (cycloalkyl)alkyl may optionally be substituted by one or more halogen atoms,
- J in each case independently of one another represent an optionally substituted 5- or 6-membered heteroaromatic ring, where the substituents independently of one another may be selected from one to three radicals W or one or more radicals R<sup>12</sup>,
- R<sup>6</sup> independently of one another represent  $-C(=E^1)R^{19}$ ,  $-LC(=E^1)R^{19}$ ,  $-C(=E^1)LR^{19}$ ,  $-LC(=E^1)LR^{19}$ ,  $-OP(=Q)(OR^{19})_2$ ,  $-SO_2LR^{18}$  or  $-LSO_2LR^{19}$ , where each  $E^1$  independently of the others represents O, S, N-R<sup>15</sup>, N-OR<sup>15</sup>, N-N(R<sup>15</sup>)<sub>2</sub>, N-S=O, N-CN or N-NO<sub>2</sub>,
- R<sup>7</sup> represents hydrogen, C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>1</sub>-C<sub>4</sub>-haloalkyl, halogen, C<sub>1</sub>-C<sub>4</sub>-alkysthio, C<sub>1</sub>-C<sub>4</sub>-alkylsulfinyl, C<sub></sub>

- alkylsulfonyl,  $C_1$ - $C_4$ -halo-alkylthio,  $C_1$ - $C_4$ -haloalkylsulfonyl, haloalkylsulfonyl,
- R<sup>9</sup> represents C<sub>1</sub>-C<sub>4</sub>-haloalkyl, C<sub>1</sub>-C<sub>4</sub>-haloalkoxy, C<sub>1</sub>-C<sub>4</sub>-haloalkylsulfinyl or halogen,
- R<sup>10</sup> represents hydrogen, C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>1</sub>-C<sub>4</sub>-haloalkyl, halogen, cyano or C<sub>1</sub>-C<sub>4</sub>-haloalkoxy,
- R<sup>11</sup> in each case independently of one another represent in each case optionally mono- to trisubstituted  $C_1$ - $C_6$ -alkylthio,  $C_1$ - $C_6$ -alkylsulfenyl,  $C_1$ - $C_6$ -haloalkythio,  $C_1$ - $C_6$ -haloalkylsulfenyl, phenylthio or phenylsulfenyl, where the substituents independently of one another may be selected from the list consisting of W,  $-S(O)_nN(R^{16})_2$ ,  $-C(=O)R^{13}$ ,  $-L(C=O)R^{14}$ ,  $-S(C=O)LR^{14}$ ,  $-C(=O)LR^{13}$ ,  $-S(O)_nNR^{13}C(=O)R^{13}$ ,  $-S(O)_nNR^{13}C(=O)R^{14}$ , and  $-S(O)_nNR^{13}S(O)_2LR^{14}$ ,
- L in each case independently of one another represent O, NR<sup>18</sup> or S,
- in each case independently of one another represent -B(OR<sup>17</sup>)<sub>2</sub>, amino, SH, thiocyanato,  $C_3$ - $C_8$ -trialkylsilyloxy,  $C_1$ - $C_4$ -alkyl disulfide, -SF<sub>5</sub>, -C(=E)R<sup>19</sup>, -LC(=E)R<sup>19</sup>, -C(=E)LR<sup>19</sup>, -LC(=E)LR<sup>19</sup>, -OP(=Q)(OR<sup>19</sup>)<sub>2</sub>, -SO<sub>2</sub>LR<sup>19</sup> or -LSO<sub>2</sub>LR<sup>19</sup>,
- Q represents O or S,
- in each case independently of one another represent hydrogen or represent in each case optionally mono- or polysubstituted C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>2</sub>-C<sub>6</sub>-alkenyl, C<sub>2</sub>-C<sub>6</sub>-alkynyl or C<sub>3</sub>-C<sub>6</sub>-cycloalkyl, where the substituents independently of one another may be selected from the group consisting of R<sup>6</sup>, halogen, cyano, nitro, hydroxyl, C<sub>1</sub>-C<sub>4</sub>-alkoxy, C<sub>1</sub>-C<sub>4</sub>-alkylsulfinyl, C<sub>1</sub>-C<sub>4</sub>-alkylsulfonyl, C<sub>1</sub>-C<sub>4</sub>-alkylamino, C<sub>2</sub>-C<sub>8</sub>-dialkylamino, C<sub>3</sub>-C<sub>6</sub>-cycloalkylamino and (C<sub>1</sub>-C<sub>4</sub>-alkyl)C<sub>3</sub>-C<sub>6</sub>-cycloalkylamino,
- R<sup>14</sup> in each case independently of one another represent in each case mono- or polysubstituted C<sub>1</sub>-C<sub>20</sub>-alkyl, C<sub>2</sub>-C<sub>20</sub>-alkenyl, C<sub>2</sub>-C<sub>20</sub>-alkynyl or C<sub>3</sub>-C<sub>6</sub>-cycloalkyl, where the substituents independently of one another may be selected from the group consisting of R<sup>6</sup>, halogen, cyano, nitro, hydroxyl, C<sub>1</sub>-C<sub>4</sub>-alkoxy, C<sub>1</sub>-C<sub>4</sub>-alkylsulfinyl, C<sub>1</sub>-C<sub>4</sub>-alkylsulfonyl,

- C<sub>1</sub>-C<sub>4</sub>-alkylamino, C<sub>2</sub>-C<sub>8</sub>-dialkylamino, C<sub>3</sub>-C<sub>6</sub>-cycloalkylamino and (C<sub>1</sub>-C<sub>4</sub>-alkyl)C<sub>3</sub>-C<sub>6</sub>-cycloalkylamino or represent optionally substituted phenyl, where the substituents independently of one another may be selected from one to three radicals W or one or more radicals R<sup>12</sup>,
- in each case independently of one another represent hydrogen or represent in each case mono- or polysubstituted C<sub>1</sub>-C<sub>6</sub>-haloalkyl or C<sub>1</sub>-C<sub>6</sub>-alkyl, where the substituents independently of one another may be selected from the group consisting of cyano, nitro, hydroxyl, C<sub>1</sub>-C<sub>4</sub>-alkoxy, C<sub>1</sub>-C<sub>4</sub>-haloalkoxy, C<sub>1</sub>-C<sub>4</sub>-alkylthio, C<sub>1</sub>-C<sub>4</sub>-alkylsulfinyl, C<sub>1</sub>-C<sub>4</sub>-alkylsulfonyl, C<sub>1</sub>-C<sub>4</sub>-haloalkylthio, C<sub>1</sub>-C<sub>4</sub>-haloalkylsulfinyl, C<sub>1</sub>-C<sub>4</sub>-haloalkylsulfonyl, C<sub>1</sub>-C<sub>4</sub>-alkylamino, C<sub>2</sub>-C<sub>8</sub>-dialkylamino, C<sub>2</sub>-C<sub>6</sub>-alkoxycarbonyl, C<sub>2</sub>-C<sub>6</sub>-alkylcarbonyl, C<sub>3</sub>-C<sub>6</sub>-trialkylsilyl and optionally substituted phenyl, where the substituents independently of one another may be selected from one to three radicals W or one or more radicals R<sup>12</sup>, or N(R<sup>15</sup>)<sub>2</sub> represents a cycle which forms the ring M,
- $R^{16}$  represents  $C_1$ - $C_{12}$ -alkyl or  $C_1$ - $C_{12}$ -haloalkyl, or  $N(R^{16})_2$  represents a cycle which forms the ring M,
- $R^{17}$  in each case independently of one another represent hydrogen or  $C_1$ - $C_4$ -alkyl, or  $B(OR^{17})_2$  represents a ring, where the two oxygen atoms are attached via a chain to two or three carbon atoms which are optionally substituted by one or two substituents independently of one another selected from the group consisting of methyl and  $C_2$ - $C_6$ -alkoxycarbonyl,
- $R^{18}$  in each case independently of one another represent hydrogen,  $C_1$ - $C_6$ -alkyl or  $C_1$ - $C_6$ -haloalkyl, or  $N(R^{13})(R^{18})$  represents a cycle which forms the ring M,
- $R^{19}$  in each case independently of one another represent hydrogen or represent in each case optionally mono- or polysubstituted  $C_1$ - $C_6$ -alkyl, where the substituents independently of one another may be selected from the group consisting of cyano, nitro, hydroxyl,  $C_1$ - $C_4$ -alkoxy,  $C_1$ - $C_4$ -haloalkylthio,  $C_1$ - $C_4$ -alkylsulfinyl,  $C_1$ - $C_4$ -alkylsulfonyl,  $C_1$ - $C_4$ -haloalkylthio,  $C_1$ - $C_4$ -haloalkylsulfinyl,  $C_1$ - $C_4$ -haloalkylsulfonyl,  $C_1$ - $C_4$ -haloalkylsulfonyl, C

alkylamino,  $C_2$ - $C_8$ -dialkylamino,  $CO_2H$ ,  $C_2$ - $C_6$ -alkoxycarbonyl,  $C_2$ - $C_6$ -alkylcarbonyl,  $C_3$ - $C_6$ -trialkylsilyl and optionally substituted phenyl, where the substituents independently of one another may be selected from one to three radicals W,  $C_1$ - $C_6$ -haloalkyl,  $C_3$ - $C_6$ -cycloalkyl or phenyl or pyridyl, each of which is optionally mono- to trisubstituted by W,

in each case represents an optionally mono- to tetrasubstituted ring which, in addition to the nitrogen atom which is attached to the substituent pair R<sup>13</sup> and R<sup>18</sup>, (R<sup>15</sup>)<sub>2</sub> or (R<sup>16</sup>)<sub>2</sub>, contains two to six carbon atoms and optionally additionally a further nitrogen, sulfur or oxygen atom, and where the substituents independently of one another may be selected from the group consisting of C<sub>1</sub>-C<sub>2</sub>-alkyl, halogen, cyano, nitro and C<sub>1</sub>-C<sub>2</sub>-alkoxy,

in each case independently of one another represent C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>2</sub>-C<sub>4</sub>-alkenyl, C<sub>2</sub>-C<sub>4</sub>-alkynyl, C<sub>3</sub>-C<sub>6</sub>-cycloalkyl, C<sub>1</sub>-C<sub>4</sub>-haloalkyl, C<sub>2</sub>-C<sub>4</sub>-haloalkenyl, C<sub>2</sub>-C<sub>4</sub>-haloalkynyl, C<sub>3</sub>-C<sub>6</sub>-halocycloalkyl, halogen, cyano, nitro, C<sub>1</sub>-C<sub>4</sub>-alkoxy, C<sub>1</sub>-C<sub>4</sub>-haloalkoxy, C<sub>1</sub>-C<sub>4</sub>-alkylthio, C<sub>1</sub>-C<sub>4</sub>-alkylsulfinyl, C<sub>1</sub>-C<sub>4</sub>-alkylsulfonyl, C<sub>1</sub>-C<sub>4</sub>-alkylamino, C<sub>2</sub>-C<sub>8</sub>-dialkylamino, C<sub>3</sub>-C<sub>6</sub>-cycloalkylamino, (C<sub>1</sub>-C<sub>4</sub>-alkylcarbonyl, C<sub>2</sub>-C<sub>6</sub>-alkoxycarbonyl, C<sub>2</sub>-C<sub>6</sub>-alkylamino-carbonyl, C<sub>3</sub>-C<sub>8</sub>-dialkylaminocarbonyl or C<sub>3</sub>-C<sub>6</sub>-trialkylsilyl,

- n in each case independently of one another represent 0 or 1,
- p in each case independently of one another represent 0, 1 or 2,

where in the case that (a)  $R^5$  represents hydrogen,  $C_1$ - $C_6$ -alkyl,  $C_1$ - $C_6$ -haloalkyl,  $C_2$ - $C_6$ -haloalkenyl,  $C_2$ - $C_6$ -haloalkynyl,  $C_1$ - $C_4$ -haloalkoxy,  $C_1$ - $C_4$ -haloalkylthio or halogen and (b)  $R^8$  represents hydrogen,  $C_1$ - $C_6$ -alkyl,  $C_1$ - $C_6$ -haloalkyl,  $C_2$ - $C_6$ -haloalkenyl,  $C_2$ - $C_6$ -haloalkynyl,  $C_1$ - $C_4$ -haloalkoxy,  $C_1$ - $C_4$ -haloalkylthio, halogen,  $C_2$ - $C_4$ -alkylcarbonyl,  $C_2$ - $C_6$ -alkoxycarbonyl,  $C_2$ - $C_6$ -alkylaminocarbonyl or  $C_3$ - $C_8$  dialkylaminocarbonyl, (c) at least one substituent selected from the group consisting of  $R^6$ ,  $R^{11}$  and  $R^{12}$  is present and (d), if  $R^{12}$  is not present, at least one  $R^6$  or  $R^{11}$  is different from  $C_2$ - $C_6$ -alkylcarbonyl,  $C_2$ - $C_6$  alkoxycarbonyl,  $C_2$ - $C_6$ -

alkylaminocarbonyl and C<sub>3</sub>-C<sub>8</sub>-dialkylaminocarbonyl, and the compounds of the general formula (I) also include N-oxides and salts,

and at least one active compound from the group of the pyrethroids (active compounds of group 2) is pyrethroid compound in a synergistically effective amount and wherein said composition is suitable for controlling animal pests.

2. (Currently amended) The composition as claimed in according to claim 1, wherein said compound of formula I is a compound comprising at least one active compound from the group of the anthranilamides of the formula (I-1) in which

$$R^3$$
 $N$ 
 $N$ 
 $R^7$ 
 $R^4$ 
 $N$ 
 $N$ 
 $N$ 
 $N$ 
 $N$ 
 $R^9$ 

in which

R<sup>2</sup> represents hydrogen or C<sub>1</sub>-C<sub>6</sub>-alkyl,

R<sup>3</sup> represents C<sub>1</sub>-C<sub>6</sub>-alkyl which is optionally substituted by a radical R<sup>6</sup>,

 $R^4$  represents  $C_1$ - $C_4$ -alkyl,  $C_1$ - $C_2$ -haloalkyl,  $C_1$ - $C_2$ -haloalkoxy or halogen,

R<sup>5</sup> represents hydrogen, C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>1</sub>-C<sub>2</sub>-haloalkyl, C<sub>1</sub>-C<sub>2</sub>-haloalkoxy or halogen,

 $R^6$  represents  $-C(=E^2)R^{19}$ ,  $-LC(=E^2)R^{19}$ ,  $-C(=E^2)LR^{19}$  or  $-LC(=E^2)LR^{19}$ , where each  $E^2$  independently of the others represents O, S, N-R<sup>15</sup>, N-OR<sup>15</sup>, N-N(R<sup>15</sup>)<sub>2</sub>, and each L independently of the others represents O or NR<sup>18</sup>,

R<sup>7</sup> represents C<sub>1</sub>-C<sub>4</sub>-haloalkyl or halogen,

 $R^9$  represents  $C_1$ - $C_2$ -haloalkyl,  $C_1$ - $C_2$ -haloalkoxy,  $S(O)_pC_1$ - $C_2$ -haloalkyl or halogen,

 $R^{15}$  in each case independently of one another represent hydrogen or represent in each case optionally substituted  $C_1$ - $C_6$ -haloalkyl or  $C_1$ - $C_6$ -alkyl, where

the substituent independently of one another may be selected from the group consisting of cyano,  $C_1$ - $C_4$ -alkoxy,  $C_1$ - $C_4$ -haloalkoxy,  $C_1$ - $C_4$ -alkyl-thio,  $C_1$ - $C_4$ -alkyl-sulfonyl,  $C_1$ - $C_4$ -haloalkylsulfinyl or  $C_1$ - $C_4$ -haloalkylsulfonyl,

R<sup>18</sup> in each case represents hydrogen or C<sub>1</sub>-C<sub>4</sub>-alkyl,

R<sup>19</sup> in each case independently of one another represent hydrogen or C<sub>1</sub>-C<sub>6</sub>-alkyl,

p independently of one another represents 0, 1, 2.

- 3. (Currently amended) The composition as claimed in according to claim 1 or 2, wherein said pyrethroid compound is comprising at least one active compound from the group of the pyrethroids (active compounds of group 2) selected from the group consisting of
  - (2-1) acrinathrin

and/or

(2-2) alpha-cypermethrin

and/or

(2-3) betacyfluthrin

## and/or

## (2-4) cyhalothrin

#### and/or

## (2-5) cypermethrin

## and/or

## (2-6) deltamethrin

#### and/or

## (2-7) esfenvalerate

### and/or

## (2-8) ethofenprox

$$H_5C_2O$$
 $H_3C$ 
 $CH_3$ 

#### and/or

## (2-9) fenpropathrin

#### and/or

## (2-10) fenvalerate

#### and/or

## (2-11) flucythrinate

#### and/or

## (2-12) lambda-cyhalothrin

#### and/or

## (2-13) permethrin

and/or

# (2-14) taufluvalinate

and/or

## (2-15) tralomethrin

$$Br_3C$$
 $CH_3$ 
 $O$ 
 $CN$ 

and/or

### (2-16) zeta-cypermethrin

and/or

## (2-17) cyfluthrin

and/or

## (2-18) bifenthrin

and/or

# (2-19) cycloprothrin

#### and/or

## (2-20) eflusilanate

#### and/or

## (2-21) fubfenprox

#### and/or

## (2-22) pyrethrin

$$R^{20} = -CH_3 \text{ or } -CO_2CH_3$$

$$R^{21} = -CH = CH_2 \text{ or } -CH_3 \text{ or } -CH_2CH_3$$

#### and/or

## (2-23) resmethrin

and/or and

(2-24) gamma-cyhalothrin

- 4. (Currently amended) The composition as claimed in claim 1, 2 or 3 comprising anthranilamides of the according to claim 1 or 2, wherein said anthranilamide of formula (I) and said at least one pyrethroid (group 2) compound are in a ratio of from 50:1 to 1:5.
- 5. (Currently amended) A method for controlling animal pests comprising contacting animal pests with The use of a synergistically effective mixture comprising compounds a compound of the formula (I) as set forth in claim 1 or 2 and said at least one pyrethroid compound. (group 2) for controlling animal pests.
- 6. (Currently amended) A process for preparing pesticides, characterized in that a synergistically effective mixture comprising compounds of the formula (I) as set forth in claim 1 or 2 and at least one pyrethroid (group 2) are mixed comprising mixing the composition according to claim 1 or 2 with extenders[[,]] and/or or surfactants[[,]] or a mixture thereof.
- 7. (New) The composition according to claim 3, wherein said anthranilamide of formula (I) and said at least one pyrethroid compound are in a ratio of from 50:1 to 1:5.